



KANK2 gene

KN motif and ankyrin repeat domains 2

Normal Function

The *KANK2* gene provides instructions for making a protein called SRC-interacting protein (SIP). SIP regulates proteins called steroid receptor coactivators (SRCs), which play critical roles in turning on (activating) certain genes. Within cells, SIP attaches (binds) to SRCs in the fluid surrounding the nucleus (the cytoplasm). By holding SRCs in the cytoplasm, SIP prevents these proteins from entering the nucleus to activate genes. In this way, SIP helps to control gene activity.

SIP is found in many organs and tissues, including the skin and kidneys. Little is known about its specific roles in various parts of the body.

Health Conditions Related to Genetic Changes

keratoderma with woolly hair

At least one mutation in the *KANK2* gene has been found to cause a form of keratoderma with woolly hair classified as type IV. It is characterized by thick, calloused skin on the palms of the hands and soles of the feet (palmoplantar keratoderma); coarse, dry, fine, tightly curled, and sparse scalp hair; and abnormal fingernails and toenails. Unlike other forms of keratoderma with woolly hair, type IV does not appear to cause heart disease.

The identified mutation changes a single protein building block (amino acid) in SIP, replacing the amino acid alanine with the amino acid valine at position 670 (written as Ala670Val or A670V). Researchers predict that this mutation affects a region of SIP that is critical for its interaction with SRCs, which would ultimately lead to changes in gene activity. However, it is unclear how this mutation affects growth of the hair and skin, or why its effects appear to be limited to these tissues.

other disorders

At least two other mutations in the *KANK2* gene have been identified in families with a form of kidney disease called nephrotic syndrome. This condition is characterized by an inability of the kidneys to filter waste products from blood, which leads to protein in the urine, swelling (edema), and possible kidney failure. Both of the *KANK2* gene mutations associated with this condition change single amino acids in SIP.

These changes are predicted to impair the function of the protein. Research shows that SIP is necessary for the normal function of kidney cells called podocytes, which


are critical for filtering waste products from the blood. However, it is unknown exactly how an altered SIP protein affects podocyte function.

It is unclear why some *KANK2* gene mutations affect the skin and hair, while others affect the kidneys. Researchers suspect that the mutations may occur in different areas of the gene and have different effects on SIP structure and function.

Chromosomal Location

Cytogenetic Location: 19p13.2, which is the short (p) arm of chromosome 19 at position 13.2

Molecular Location: base pairs 11,164,267 to 11,197,567 on chromosome 19 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- ANKRD25
- ankyrin repeat domain 25
- ankyrin repeat domain-containing protein 25
- KIAA1518
- kidney ankyrin repeat-containing protein 2
- matrix-remodeling-associated protein 3
- matrix-remodelling associated 3
- MXRA3
- PPKWH
- SIP
- SRC-1 interacting protein
- SRC-1-interacting protein

- SRC-interacting protein
- SRC1-interacting protein

Additional Information & Resources

Scientific Articles on PubMed

- PubMed
https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28KANK2%5BTIAB%5D%29+OR+%28KN+motif+and+ankyrin+repeat+domains+2%5BTIAB%5D%29%29+OR+%28SRC-interacting+protein%5BTIAB%5D%29+OR+%28%28SRC*%5BTIAB%5D%29+AND+%28SIP%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D

OMIM

- KN MOTIF- AND ANKYRIN REPEAT DOMAIN-CONTAINING PROTEIN 2
<http://omim.org/entry/614610>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_KANK2.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=KANK2%5Bgene%5D>
- HGNC Gene Family: Ankyrin repeat domain containing
<http://www.genenames.org/cgi-bin/genefamilies/set/403>
- HGNC Gene Family: KN motif and ankyrin repeat domain containing
<http://www.genenames.org/cgi-bin/genefamilies/set/623>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=29300
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/25959>
- UniProt
<http://www.uniprot.org/uniprot/Q63ZY3>

Sources for This Summary

- Gee HY, Zhang F, Ashraf S, Kohl S, Sadowski CE, Vega-Warner V, Zhou W, Lovric S, Fang H, Nettleton M, Zhu JY, Hoefele J, Weber LT, Podracka L, Boor A, Fehrenbach H, Innis JW, Washburn J, Levy S, Lifton RP, Otto EA, Han Z, Hildebrandt F. KANK deficiency leads to podocyte dysfunction and nephrotic syndrome. *J Clin Invest*. 2015 Jun;125(6):2375-84. doi: 10.1172/JCI79504. Epub 2015 May 11.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/25961457>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4497755/>
- OMIM: KN MOTIF- AND ANKYRIN REPEAT DOMAIN-CONTAINING PROTEIN 2
<http://omim.org/entry/614610>
- Ramot Y, Molho-Pessach V, Meir T, Alper-Pinus R, Siam I, Tams S, Babay S, Zlotogorski A. Mutation in KANK2, encoding a sequestering protein for steroid receptor coactivators, causes keratoderma and woolly hair. *J Med Genet*. 2014 Jun;51(6):388-94. doi: 10.1136/jmedgenet-2014-102346. Epub 2014 Mar 26.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/24671081>
- Zhang Y, Zhang H, Liang J, Yu W, Shang Y. SIP, a novel ankyrin repeat containing protein, sequesters steroid receptor coactivators in the cytoplasm. *EMBO J*. 2007 Jun 6;26(11):2645-57. Epub 2007 May 3.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17476305>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1888672/>
- Zhu Y, Kakinuma N, Wang Y, Kiyama R. Kank proteins: a new family of ankyrin-repeat domain-containing proteins. *Biochim Biophys Acta*. 2008 Feb;1780(2):128-33. Epub 2007 Oct 4.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17996375>

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/gene/KANK2>

Reviewed: November 2015

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services